

SUMMATIVE ASSESSMENT - II - 2016 - 2017

CLASS-X - MATHS - PAPER-I

Part - A & B

KEY

Class : X

Part - A

Marks : 60

Section - I (Each question carries 1 mark)

$$1. \quad \left. \begin{aligned} A &= \{x : x \text{ is a prime factor of } 30\} = \{2, 3, 5\} \\ B &= \{x : x \text{ is a prime below of } 20\} = \{2, 3, 5, 7, 11, 13, 17, 19\} \end{aligned} \right\} \frac{1}{2}m$$

$$\begin{aligned} \text{(i)} \quad A \cup B &= \{2, 3, 5\} \cup \{2, 3, 5, 7, 11, 13, 17, 19\} \\ &= \{2, 3, 5, 7, 11, 13, 17, 19\} \\ &= B \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad A \cap B &= \{2, 3, 5\} \cap \{2, 3, 5, 7, 11, 13, 17, 19\} \\ &= \{2, 3, 5\} \\ &= A \end{aligned}$$

1m

$$2. \quad \text{Given Q.E } 3x^2 - 2x + \frac{1}{3} = 0$$

$$\text{Here } a = 3, b = -2, c = \frac{1}{3}$$


$$\begin{aligned} \Delta &= b^2 - 4ac \\ &= (-2)^2 - 4(3)\left(\frac{1}{3}\right) \end{aligned} \left\} \frac{1}{2}$$

$$= 4 - 4$$

$$= 0 \quad \therefore \text{Roots are real and equal } -\frac{1}{2}$$


1m

3. If $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$ are a pair of linear equations and if $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$ then the system of linear equations are in 'consistent' system. 1m



(or) For any such figure $\left. \begin{aligned} \text{Figure } -\frac{1}{2} \\ \text{Name } -\frac{1}{2} \end{aligned} \right\} 1m$

7.



Radius of the sphere = $6\sqrt{3}$ cm

when largest possible 'cube' is carved out of sphere, then diagonal of the cube = diameter of sphere $\left. \vphantom{\begin{array}{l} \text{when largest possible 'cube' is carved out of sphere,} \\ \text{then diagonal of the cube = diameter of sphere} \end{array}} \right\} \frac{1}{2}$

$$\begin{aligned} \therefore d &= 2(6\sqrt{3}) \\ \sqrt{3} S &= 12\sqrt{3} \end{aligned} \left. \vphantom{\begin{array}{l} \therefore d = 2(6\sqrt{3}) \\ \sqrt{3} S = 12\sqrt{3} \end{array}} \right\} \frac{1}{2}$$

$$S = 12$$

$$\begin{aligned} \therefore \text{Surface area of cube} &= 6S^2 \\ &= 6(12)^2 = 864 \text{ cm}^2 \end{aligned} \left. \vphantom{\begin{array}{l} \therefore \text{Surface area of cube} = 6S^2 \\ = 6(12)^2 = 864 \text{ cm}^2 \end{array}} \right\} 1 \text{ cm} \quad 2\text{m}$$

8. Assume that $\frac{1}{3\sqrt{2}}$ is rational

$$\text{Let } \frac{1}{3\sqrt{2}} = \frac{p}{q} \text{ (q} \neq 0, p, q \text{ are co-primes)} \left. \vphantom{\text{Let } \frac{1}{3\sqrt{2}} = \frac{p}{q}} \right\} \frac{1}{2}$$

$$\begin{aligned} 3\sqrt{2}p &= q \\ \sqrt{2} &= \frac{q}{3p} \end{aligned} \left. \vphantom{\begin{array}{l} 3\sqrt{2}p = q \\ \sqrt{2} = \frac{q}{3p} \end{array}} \right\} \frac{1}{2}$$

$$\text{Here LHS is an irrational and } \frac{q}{3p} \text{ is rational.} \left. \vphantom{\text{Here LHS is an irrational and } \frac{q}{3p} \text{ is rational.}} \right\} \frac{1}{2}$$

This is a contradiction

$$\begin{aligned} \therefore \text{Our assumption is false.} \\ \therefore \frac{1}{3\sqrt{2}} \text{ is irrational.} \end{aligned} \left. \vphantom{\begin{array}{l} \therefore \text{Our assumption is false.} \\ \therefore \frac{1}{3\sqrt{2}} \text{ is irrational.} \end{array}} \right\} \frac{1}{2} \quad 2\text{m}$$

9. Let no. of honey bees = x
no. of flowers = y $\left. \vphantom{\begin{array}{l} \text{Let no. of honey bees = x} \\ \text{no. of flowers = y} \end{array}} \right\} \frac{1}{2}$

(i) Two honey bees sit on each flower, one bee was left out

$$\begin{aligned} \therefore x &= 2y + 1 \\ \Rightarrow x - 2y &= 1 \text{ (1)} \end{aligned} \left. \vphantom{\begin{array}{l} \therefore x = 2y + 1 \\ \Rightarrow x - 2y = 1 \end{array}} \right\} \frac{1}{2}$$

(ii) Three bees sit on each flower, no flower is left

$$\begin{aligned} \therefore y &= \frac{x}{3} + 0 \\ \Rightarrow x - 3y &= 0 \text{ (2)} \end{aligned} \left. \vphantom{\begin{array}{l} \therefore y = \frac{x}{3} + 0 \\ \Rightarrow x - 3y = 0 \end{array}} \right\} \frac{1}{2} \quad 2\text{m}$$

Section - III (Each question carries 4 marks)

10 (a) Given Q.E $x^2 - (m+1)x + 6 = 0$

$$\begin{aligned} x = 3 &\Rightarrow (3)^2 - (m+1)3 + 6 = 0 \\ 9 - (3m+3) + 6 &= 0 \\ -3m + 12 &= 0 \\ m &= 4 \end{aligned} \quad \left. \begin{array}{l} \\ \\ \\ \end{array} \right\} \begin{array}{l} 1 \\ 1 \end{array}$$

Now the equation become


$$\begin{aligned} x^2 - (4+1)x + 6 &= 0 \\ x^2 - 5x + 6 &= 0 \end{aligned} \quad \left. \begin{array}{l} \\ \end{array} \right\} 1$$

$$(x-2)(x-3) = 0 \Rightarrow x = \{2, 3\} \quad \left. \begin{array}{l} \\ \end{array} \right\} 1$$

\therefore Other root = 2

4m

10 (b)



Diameter of sphere = 28 cm,

$$\begin{aligned} \text{Diameter of cone} &= 4\frac{2}{3} \text{ cm} \\ \text{radius} &= \frac{14}{3} \times \frac{1}{2} = \frac{7}{3} \text{ cm} \end{aligned} \quad \left. \begin{array}{l} \\ \end{array} \right\} 1$$

$$r = \frac{28}{2} = 14 \text{ cm} \dots\dots\dots (1) \quad \left. \begin{array}{l} \\ \end{array} \right\} 1$$

height = 3cm

ATP

$$\begin{aligned} \text{The sphere is melted and cost into some (n) cones} \\ \therefore n \times \text{volume of each cone} &= \text{volume of sphere} \end{aligned} \quad \left. \begin{array}{l} \\ \end{array} \right\} \frac{1}{2}$$

$$n \times \frac{1}{3} \pi r^2 h = \frac{4}{3} \pi r^3$$

$$n \times \frac{1}{3} \times \frac{22}{7} \times \frac{7}{3} \times \frac{7}{3} \times 3 = \frac{2}{3} \times \frac{22}{7} \times 14 \times 14 \times 14$$

$$\therefore n = 672 \quad \left. \begin{array}{l} \\ \end{array} \right\} \frac{1}{2}$$

4m

11 (a) $x^2 + y^2 = 6xy$

Add '2xy' on both sides

$$\begin{aligned} x^2 + y^2 + 2xy &= 6xy + 2xy \\ (x+y)^2 &= 8xy \end{aligned} \quad \left. \begin{array}{l} \\ \end{array} \right\} 1$$

Apply 'log' on both sides

$$\log (x+y)^2 = \log 8xy \quad \left. \begin{array}{l} \\ \end{array} \right\} 1$$

$$\begin{aligned} 2 \log (x+y) &= \log 8 + \log x + \log y \\ &= \log 2^3 + \log x + \log y \\ &= 3 \log 2 + \log x + \log y \end{aligned} \quad \left. \begin{array}{l} \\ \\ \end{array} \right\} \begin{array}{l} 1 \quad [\because \log xy = \log x + \log y] \\ 1 \quad [\because \log x^m = m \log x] \end{array}$$

Hence proved

4m

(b) Let 'a' be a positive integer

Take $b = 3$, Acc to Euclids division Lemma $\left. \begin{array}{l} a = bq + r, r = 0, 1, 2 (\because 0 \leq r < b) \end{array} \right\} \frac{1}{2}$

$$\Rightarrow \left. \begin{array}{l} a = 3q \\ = 3q + 1 \\ = 3q + 2 \end{array} \right\} \frac{1}{2}$$

(i) If $a = 3q$

$$\Rightarrow \left. \begin{array}{l} a^3 = (3q)^3 \\ = 27q^3 \\ = 9(3q^3) = 9m \end{array} \right\} 1$$

(ii) $a = 3q + 1$

$$\Rightarrow \left. \begin{array}{l} a^3 = (3q+1)^3 \\ = 27q^3 + 27q^2 + 9q + 1 \\ = 9(3q^3 + 3q^2 + q) + 1 \\ = 9m + 1 \end{array} \right\} 1$$

(iii) If $a = 3q + 2$

$$\Rightarrow \left. \begin{array}{l} a^3 = (3q+2)^3 \\ = 27q^3 + 54q^2 + 36q + 8 \\ = 9(3q^3 + 6q^2 + 4q) + 8 \\ = 9m + 8 \end{array} \right\} 1$$

For any positive integer, the cube is always in the form of

$9m, 9m + 1$ or $9m+8$

Hence proved

4m

12 (a) Let the speed of the car = x kmph

distance travelled = 36 km

$$\therefore \text{time taken} = \frac{36}{x} \dots\dots\dots (1)$$

If speed is increased by 10 kmph, then $\left. \begin{array}{l} \end{array} \right\} 1$

$$\text{time taken} = \frac{36}{x+10} \dots\dots\dots (2)$$

ATP

$$\text{Difference in time taken} = 18 \text{ min}$$

$$= \frac{18}{60} \text{ hr} \left. \begin{array}{l} \end{array} \right\} 1$$


$$\therefore \frac{36}{x} - \frac{36}{x+10} = \frac{18}{60}$$

$$\text{Simplifying } x^2 + 10x - 1200 = 0 \left. \begin{array}{l} \end{array} \right\} 1$$

$$\begin{array}{l} \text{Solving } (x+40)(x-30) = 0 \\ \Rightarrow x = \{-40, 30\} \end{array} \left. \begin{array}{l} \end{array} \right\} 1$$

Speed of the car = 30 kmph

4m



Length of the capsule = 14mm

Width of the capsule = 5mm

The capsule is the combination of two hemispheres and a cylinder

$$\therefore \text{radius of hemisphere} = \frac{d}{2} = \frac{5}{2} \text{ mm}$$

$$\begin{aligned} \therefore \text{Volume of two hemispherical ends} &= 2 \times \frac{2}{3} \pi r^3 \\ &= 2 \times \frac{2}{3} \times \frac{22}{7} \times \frac{5}{2} \times \frac{5}{2} \times \frac{5}{2} \left. \vphantom{\frac{22}{7}} \right\} 1 \\ &= \frac{1375}{21} \text{ mm}^3 \end{aligned}$$

$$\begin{aligned} \text{Volume of cylinder} &= \pi r^2 h \\ &= \frac{22}{7} \times \frac{5}{2} \times \frac{5}{2} \times \left(14 - 2 \left(\frac{5}{2} \right) \right) \left. \vphantom{\frac{22}{7}} \right\} 1 \\ &= \frac{22}{7} \times \frac{5}{2} \times \frac{5}{2} \times 9 \text{ mm}^3 = \frac{2475}{14} \text{ mm}^3 \end{aligned}$$

\therefore Total volume of capsule

$$\begin{aligned} &= \text{Volume of two hemispherical ends} + \text{Volume of cylinder} \\ &= \frac{1375}{21} + \frac{2475}{14} \left. \vphantom{\frac{1375}{21}} \right\} 1 \\ &= 65.47 + 176.78 \left. \vphantom{\frac{1375}{21}} \right\} 1 \\ &= 242.25 \text{ mm}^3 \end{aligned}$$

4m


13 (a) For writing the table for $p(x) = x^2 - 4x - 5$ $\left. \vphantom{p(x)} \right\} 1 \frac{1}{2}$

x	-3	-2	-1	0	1	2	3	4
$p(x) = x^2 - 4x - 5$	$(-3)^2 - 4(-3) - 5$	$(-2)^2 - 4(-2) - 5$	$(-1)^2 - 4(-1) - 5$	$0^2 - 4(0) - 5$	$1^2 - 4(1) - 5$	$2^2 - 4(2) - 5$	$3^2 - 4(3) - 5$	$4^2 - 4(4) - 5$
y	16	7	0	-5	-8	-9	-8	-5
(x, y)	(-3, 16)	(-2, 7)	(-1, 0)	(0, -5)	(1, -8)	(2, -9)	(3, -8)	(4, -5)

Scale = X-axis = 1 cm = 1 unit

Y-axis = 1 cm = 2 unit

For Graph :



Check : $P(x) = x^2 - 4x - 5$

$P(-1) = (-1)^2 - 4(-1) - 5$

$= 1 + 4 - 5$

$= 0$

$P(5) = (5)^2 - 4(5) - 5$

$= 25 - 20 - 5$

$= 0$

} $1\frac{1}{2}$

- Solution : 1) The Graph of the polyminal is a curve
- 2) It cuts x-axis at $(-1, 0)$ and $(5, 0)$
- 3) Zeroes = $\{-1, 5\}$
- } $\frac{1}{2}$

4m

13 (b) For writing tables for equations

(i) $2x - y = 5$

x	0	$\frac{5}{2}$	2
y	-5	0	-1
(x, y)	(0, -5)	$(\frac{5}{2}, 0)$	(2, -1)


(ii) $4x - 2y = 12$

x	0	3	-2
y	-6	0	-10
(x, y)	(0, -6)	(3, 0)	(-2, -10)

} 2

Scale = X-axis = 1 cm = 1 unit

Y-axis = 1 cm = 2 unit



Solution : 1) The Graph of pair of linear equations is a pair of parallel lines. $\left. \begin{array}{l} \text{2) Hence the Equations are in inconsistent system.} \end{array} \right\} \frac{1}{2} \quad 4m$

SUMMATIVE ASSESSMENT - II - 2016 - 2017

CLASS-X - MATHS - PAPER-I

KEY

Class : X

Part - B

Marks : 20

III.

- 14. C
- 15. B
- 16. C
- 17. D
- 18. B
- 19. D
- 20. B
- 21. C
- 22. C
- 23. A
- 24. D
- 25. D
- 26. B
- 27. B
- 28. C
- 29. C
- 30. D
- 31. C
- 32. C
- 33. B